

# Mepilex® Ag

## Absorbent antimicrobial foam dressing

### Polyurethane foam pad

- Absorbs exudate
- Conformable
- May be used under compression


### Polyurethane backing film

- Breathable
- Shower-proof<sup>1</sup>
- Bacteria and viral barrier<sup>1-3,†</sup>

† Microbes larger than 25 nm

### Powerful silver performance

- Antimicrobial activity within 30 minutes and sustained effect for up to seven days<sup>1,5-7</sup>



### Safetac® technology layer

- Less pain during dressing changes<sup>8</sup>
- Does not stick to the wound, for less disturbance<sup>9</sup>
- Seals wound margins to avoid maceration<sup>10</sup>



## How Mepilex Ag works

Mepilex Ag is a soft and highly conformable antimicrobial foam dressing that absorbs exudate and maintains a moist wound environment. The Safetac technology layer seals the wound edges, preventing the exudate from leaking onto the surrounding skin, thus minimising the risk of maceration. The Safetac technology layer ensures that the dressing can be changed without damaging the wound or surrounding skin or exposing the patient to additional pain.

Mepilex Ag starts to inactivate wound related pathogens within 30 minutes with sustained effect for up to seven days<sup>1,5-7</sup>. By reducing the number of microorganisms, Mepilex Ag may also reduce odour.

## Benefits of Mepilex Ag

- Minimises pain and trauma at dressing changes<sup>8,9</sup>
- Stays in place allowing easy application of compression or retention bandages
- Well-suited for use under compression bandages
- May be cut to suit various wound shapes and difficult-to-dress locations
- Promotes patient comfort during wear
- May remain in place for several days depending on the condition of the wound
- May be adjusted without losing its adherent properties
- Low potential for skin irritation and allergy

## Areas of use

- Moderately exuding wounds
- Leg and foot ulcers
- Pressure injuries
- Under compression
- Partial thickness burns

**Precautions:** 1. Do not use on patients with a known sensitivity to silver. 2. Do not use Mepilex Ag or Mepilex Heel Ag during radiation treatment or examinations e.g. X-ray, ultrasound, or diathermy. 3. Do not use Mepilex Ag or Mepilex Heel Ag together with oxidising agents, such as hypochlorite solutions or hydrogen peroxide.

## Mepilex Ag ordering information†

Product Code	Size	Pcs/box
287110	10 x 10 cm	5